


Medicine in Evolution



TRANSLATIONAL AND EXPERIMENTAL CLINICAL RESEARCH CENTER IN ORAL HEALTH (TEXC-OH)
CENTER FOR CONTINUOUS MEDICAL EDUCATION

Volume XXXII, No. 2, 2026, Timișoara, Romania
ISSN 2065-376X

MEDICINE IN EVOLUTION



**TRANSLATIONAL AND EXPERIMENTAL CLINICAL
RESEARCH CENTRE IN ORAL HEALTH**

“VICTOR BABEȘ” UNIVERSITY OF MEDICINE AND PHARMACY TIMIȘOARA

<https://medicineinevolution.ro/>

**Translational and Experimental Clinical Research Center
in Oral Health**

<https://www.umft.ro/en/research-centres/ccextso/>

INDEX  COPERNICUS
I N T E R N A T I O N A L




EDITORIAL BOARD

FOUNDING EDITOR

Prof. Ancusa Mircea
MD, PhD

Prof. Podariu Angela Codruța
DMD, PhD, Timișoara



ASSOCIATE EDITORS	EDITOR IN CHIEF	ASSISTANT EDITOR
<p>Prof. Dehelean Cristina MD, PhD, Timișoara</p> <p>Prof. Oancea Cristian MD, PhD, Timișoara</p> <p>Prof. Păunescu Virgil MD, PhD, Timișoara</p> <p>Assoc. Prof. Sava-Rosianu Ruxandra DMD, PhD, Timișoara</p>	<p>Prof. Jumanca Daniela DMD, PhD, Timișoara</p> <p>Prof. Galuscan Atena DMD, PhD, Timișoara</p> <p>Prof. Oancea Roxana MD, PhD, Timișoara</p>	<p>Mădălina-Victoria Coccoceanu INF. EC., Timișoara</p>

NATIONAL EDITORIAL BOARD		
<p>Prof. Ardelean Lavinia DMD, PhD, Timișoara</p> <p>Lecturer Balean Octavia - Iulia DMD, PhD, Timișoara</p> <p>Prof. Bechir Anamaria DMD, PhD, București</p> <p>Assoc. Prof. Beresescu Liana DMD, PhD, Târgu-Mureș</p> <p>Prof. Bica Cristina Ioana Tg. Mureș</p> <p>Prof. Borza Claudia MD, PhD, Timișoara</p> <p>Prof. Bratu Emanuel DMD, PhD, Timișoara</p> <p>Dr. Brehar-Cioflec Dana MD, PhD, Timișoara</p> <p>Assoc. Prof. Bucur Adina MD, PhD, Timișoara</p> <p>Prof. Bunu Panaitescu Carmen MD, PhD, Timișoara</p> <p>Prof. Buzatu Roxana DMD, PhD, Timișoara</p> <p>Prof. Caraiane Aureliana DMD, PhD, Constanța</p>	<p>Assist. Prof. Caramida Mariana DMD, PhD, București</p> <p>Assoc. Prof. Daguci Constantin DMD, PhD, Craiova</p> <p>Prof. Dehelean Cristina Adriana PhD, Timișoara</p> <p>Assoc. Prof. Draghici George Andrei PhD, Timișoara</p> <p>Prof. Dumitrașcu Victor MD, PhD, Timișoara</p> <p>Prof. Dumitrache Adina DMD, PhD, București</p> <p>Assoc. Prof. Esian Daniela Elena DMD, PhD, Târgu-Mureș</p> <p>Prof. Forna Norina Consuela DMD, PhD, Iași</p> <p>Assoc. Prof. Funieru Cristian DMD, PhD, București</p> <p>Prof. Gălușcan Atena DMD, PhD, Timișoara</p> <p>Assist. Prof. Goția Laura DMD, PhD, Timișoara</p> <p>Prof. Jivănescu Anca DMD, PhD, Timișoara</p>	<p>Prof. Leretter Marius MD, PhD, Timișoara</p> <p>Prof. Lucaciu Ondine Patricia Cluj Napoca</p> <p>Prof. Marian Catalin Valer MD, PhD, Timisoara</p> <p>Lecturer Matichescu Anamaria DMD, PhD, Timișoara</p> <p>Assoc. Prof. Mesaros Anca Stefania DMD, PhD, Cluj-Napoca</p> <p>Prof. Mercuț Veronica DMD, PhD, Craiova</p> <p>Assoc. Prof. Muntean Alexandrina DMD, PhD, Cluj-Napoca</p> <p>Prof. Murariu Alice DMD, PhD, Iasi</p> <p>Prof. Negrutiu Meda Lavinia DMD, PhD, Timișoara</p> <p>Prof. Oancea Roxana DMD, PhD, Timișoara</p> <p>Prof. Păcurar Mariana DMD, PhD, Târgu-Mureș</p>


NATIONAL EDITORIAL BOARD

Prof. Petrescu Emanuela DMD, PhD, Timișoara	Prof. Romînu Mihai DMD, PhD, Timișoara	Prof. Șoica Codruța-Mariana PhD, Timișoara
Prof. Pinzaru Iulia Andreea PhD, Timișoara	Prof. Rusu Darian DMD, PhD, Timișoara	Prof. Stratul Stefan-Ioan MD, PhD, Timișoara
Prof. Popescu Roxana MD, PhD, Timișoara	Prof. Rusu Laura Cristina DMD, PhD, Timișoara	Prof. Székely Melinda DMD, PhD, Târgu-Mureș
Prof. Popovici Ramona Amina DMD, PhD, Timișoara	Assoc. Prof. Sava-Roșianu Ruxandra DMD, PhD, Timișoara	Assoc. Prof. Tanase Doina Alina PhD, Timișoara
Prof. Popșor Sorin DMD, PhD, Târgu Mureș	Prof. Saveanu Iulia Catalina DMD, PhD, Iasi	Prof. Tudor Liana DMD, PhD, Oradea
Prof. Porojan Liliana DMD, PhD, Timișoara	Prof. Sfeatcu Ruxandra DMD, PhD, București	Assoc. Prof. Vernic Corina MD, PhD, Timișoara
Prof. Pricop Marius DMD, PhD, Timișoara	Prof. Sinescu Cosmin DMD, PhD, Timișoara	Prof. Zetu Irina DMD, PhD, Iași

INTERNATIONAL EDITORIAL BOARD

Assist. Prof. Dr. Ambarkova Vesna Macedonia	Prof. Dr. Lingström Peter Sweden	Prof. Puriene Alina Lithuania
Prof. Campus Guglielmo Giuseppe Sweden	Prof. Nikolovska Julijana Macedonia	Prof. Soares Henrique Luis Portugal
Assoc. Prof. Dr. Getova Biljana Macedonia	Prof. Paganelli Corrado Italy	Prof. Veltri Nicola Italy
Assoc. Prof. Giraudeau Nicolas France	Assoc. Prof. Porosencova Tatiana Republic of Moldova	Lecturer Vukovic Ana Serbia
Assoc. Prof. Hysi Dorjan Albania	Prof. Plesh Octavia USA	Prof. Zimmer Stefan Germany

Background



The current journal was established by Prof. Dr. Mircea Ancusa in 1999, with the aim of acquiring knowledge and sharing insights in the noble profession guided by the principle "primum non nocere" (first, do no harm). In 2005, it was entrusted to a group of dedicated researchers at the Center of Health Education and Motivation for Prevention in Dentistry, under the leadership of Prof. Angela Codruta Podariu, DMD, PhD, at the Department of Preventive Dentistry of the University of Medicine and Pharmacy "Victor Babes" in Timisoara, Romania.

The inception of the journal stemmed from a dedication to exchange experiences in both professional and research domains. It was envisioned to encompass all medical specialties, with the aspiration that the published manuscripts would exhibit exceptional quality, elevating the journal's reputation. Esteemed professionals were enlisted to the editorial board and the review committee, individuals recognized for their expertise in the realm of research. The decision to publish papers in English was made to broaden accessibility to the global research community and enhance international recognition.

Since then, the journal has been regularly published under the auspices of the Center of Health Education and Motivation for Prevention in Dentistry, disseminating national and international research studies with the objective of evolving into a comprehensive evidence-based publication. Presently, the journal has transitioned to the stewardship of the Translational and Experimental Clinical Research Centre in Oral Health, situated within the Department of Preventive, Community Dentistry, and Oral Health. Its objectives are aligned with the vision of esteemed organizations such as the World Health Organization and the International Dental Federation, seamlessly integrating into the research strategy of Victor Babes University of Medicine and Pharmacy Timisoara.

"Medicine in Evolution" stands as a distinguished, peer-reviewed, open access journal dedicated to the dissemination of original theoretical research spanning the interdisciplinary spectrum of medicine and healthcare. Encompassing various topics within the realms of human life sciences, medical community, dental medicine, and pharmacology, the journal warmly welcomes original research papers, communications, letters, short notes, case reports, and reviews for submission. Committed to conducting rigorous peer reviews and expediting the publication of groundbreaking research, its mission is to advance the field of medicine through scholarly discourse.

CONTENTS

ARTICLES



Andrei-Bogdan Faur, Isabela Toser, Raul Nicolae Rotar, Anca-Elena Anghel-Lorinți, Anca Jivănescu

Fully Digital Chairside Workflow for Anterior Prosthetic Rehabilitation: A Clinical Case Report from Digital Data Acquisition to Final Smile Enhancement 71

Cristian Argyelan, Angelica Moldovan, Paul Morari, Ersilia Alexa

Hypoglycemic Bread as a Nutritional Strategy for the Management of Diabetes Mellitus 83

Fully Digital Chairside Workflow for Anterior Prosthetic Rehabilitation: A Clinical Case Report from Digital Data Acquisition to Final Smile Enhancement



<https://doi.org/10.70921/medev.v32i2.2032>

Andrei-Bogdan Faur^{1,2}, Isabela Toser^{1*}, Raul Nicolae Rotar^{1,2}, Anca-Elena Anghel-Lorinți¹, Anca Jivănescu^{1,2}

¹Department of Prosthodontics, University of Medicine and Pharmacy "Victor Babes", B-dul Revolutiei 1989, No. 9, 300580 Timisoara, Romania; andrei.faur@umft.ro (A.-B.F.); isabela.toser@umft.ro (I.T.); anca.anghel@umft.ro (A.-E.A.-L.); rotar.raul@umft.ro (R.R.); jivanescu.anca@umft.ro (A.J.)

²TADERP Research Center, Department of Prosthodontics, University of Medicine and Pharmacy "Victor Babes", B-dul Revolutiei 1989, No. 9, 300580 Timisoara, Romania

Correspondence to:

Name: Isabela Toser

E-mail address: isabela.toser@umft.ro

Received: 6 April 2026; Accepted: 4 May 2026; Published:

Abstract

1. Background/Objectives: The integration of intraoral scanning, CAD/CAM technologies, and additive manufacturing has enabled the development of fully digital workflows in prosthodontics; however, their application in chairside anterior rehabilitations remains less documented. This clinical case aimed to evaluate a complete digital, chairside protocol for anterior maxillary rehabilitation, emphasizing digital planning and staged provisionalization. 2. Case Presentation: A 50-year-old male patient with partial anterior maxillary edentulism was treated using a fully digital workflow, including intraoral and facial scanning, Digital Smile Design, CAD-based wax-up, and fabrication of 3D-printed eggshell and on-abutment provisional restorations, followed by chairside milling of monolithic zirconia fixed prostheses. 3. Results: The staged provisional approach enabled validation of aesthetics and function, guided minimally invasive tooth preparation, and facilitated soft tissue conditioning. The final restorations demonstrated satisfactory marginal adaptation, occlusion, and aesthetic integration, with high patient acceptance. 4. Conclusion: A fully digital, chairside workflow allowed predictable and efficient anterior rehabilitation, improving treatment control, communication, and patient involvement, supporting its clinical applicability in modern prosthodontic practice within the limitations of a single case.

Keywords: Digital dentistry; intraoral scanning; CAD/CAM; chairside workflow; anterior prosthetic rehabilitation; Digital Smile Design; 3D printing; provisional restorations; ovate pontic; monolithic zirconia.

INTRODUCTION

The integration of digital technologies into prosthodontics has fundamentally transformed the diagnosis, planning, and execution of restorative treatments. In particular, the development of intraoral scanners, computer-aided design/computer-aided manufacturing (CAD/CAM) systems, and additive manufacturing has enabled the implementation of fully digital workflows, offering new possibilities for precision, efficiency, and patient-centered care [1].

Intraoral scanning represents a key component of this transformation, providing accurate digital impressions while eliminating many of the limitations associated with conventional impression techniques. Clinical studies and systematic reviews have demonstrated that intraoral scanners can achieve comparable or superior accuracy in fixed prosthodontics, while also improving patient comfort and reducing clinical time [2,3]. These advantages are especially relevant in anterior rehabilitations, where high esthetic demands require precise data acquisition and reproducibility.

Beyond data acquisition, the integration of digital tools into treatment planning has enabled the concept of facially driven prosthetic design, where dental restorations are planned in harmony with facial features and smile dynamics. Digital Smile Design (DSD) has emerged as a valuable tool for improving communication between clinician, technician, and patient, as well as for enhancing predictability by allowing visualization of the final outcome prior to treatment initiation [4].

The evolution toward a fully digital chairside workflow from intraoral scanning to CAD design and CAM fabrication has further streamlined clinical and laboratory procedures. Such workflows can reduce manual errors, improve consistency between treatment phases, and allow rapid iteration of designs, particularly when combined with additive manufacturing technologies such as 3D printing [1,5]. The use of 3D-printed provisional restorations has gained increasing attention due to their cost-efficiency, reproducibility, and ability to accurately transfer digital designs into the clinical environment [6].

Provisional restorations play a critical role in prosthodontic treatment, particularly in the anterior region, where they serve not only as temporary restorations but also as diagnostic tools for evaluating esthetics, function, and soft tissue response. Concepts such as ovate pontic design and staged provisionalization have been shown to support gingival contouring and emergence profile development, contributing to improved esthetic outcomes [7]. When integrated within a digital workflow, provisional restorations can be precisely designed and modified, allowing progressive validation of the treatment plan before fabrication of definitive restorations.

Additionally, the development of chairside CAD/CAM systems has enabled clinicians to perform multiple steps of the restorative process within a single clinical setting. Chairside workflows offer advantages such as reduced treatment time, elimination of conventional laboratory steps, and increased control over the entire process, although they may be limited by equipment capabilities and material constraints [8].

Despite these advancements, the application of fully digital, chairside workflows in multi-unit, tooth-supported anterior rehabilitations remain less documented compared to single-unit or implant-supported restorations. Therefore, further clinical reports are necessary to evaluate their feasibility, advantages, and limitations.

Aim and objectives

The aim of this clinical case report is to describe a complete digital chairside workflow for anterior maxillary prosthetic rehabilitation, highlighting the role of intraoral scanning,

facially driven design, staged provisionalization, and CAD/CAM fabrication in achieving a predictable functional and esthetic outcome.

CASE DESCRIPTION

Ethics, Case Introduction, and Clinical Objectives

This clinical case was approved by the Research Ethics Committee of the Victor Babeş University of Medicine and Pharmacy (Approval No. 62/06.11.2025) and was conducted in accordance with the principles outlined in the Declaration of Helsinki. The patient provided written informed consent for participation, including permission for the use of clinical data and photo-video documentation for scientific purposes.

A 50-year-old male patient presented to the Department of Prosthodontics, Faculty of Dental Medicine, seeking prosthetic rehabilitation for missing maxillary right canine and maxillary left lateral incisor (teeth 1.3 and 2.2). The patient was informed about all available treatment options, including implant-supported prosthetic rehabilitation; however, he declined implant therapy and opted for a fixed dental prosthetic solution supported by natural abutment teeth.

Clinical and digital evaluation protocols were implemented, including intraoral examination, photographic documentation, intraoral and facial scanning, and radiographic assessment. Based on these findings, a fully digital chairside workflow was selected.

Initial Clinical Examination and Digital Diagnostic Protocol

A comprehensive intraoral examination was performed, revealing no clinical signs of periodontal disease. The periodontal tissues were clinically healthy, with no evidence of gingival inflammation, bleeding on probing, attachment loss, or pathological tooth mobility. The oral hygiene status was satisfactory, and the remaining dentition exhibited stable periodontal support, making it suitable for prosthetic rehabilitation using natural abutments.

Occlusal analysis identified the presence of premature contacts and minor occlusal interferences, likely associated with the loss of dental units and compensatory tooth wear. These discrepancies were managed through selective enamel recontouring, aiming to establish a stable and functional occlusal scheme prior to initiating the digital diagnostic workflow. This preliminary equilibration step was essential to ensure an accurate and reproducible maxillomandibular relationship during intraoral scanning and digital bite registration.

An evaluation of the vertical dimension of occlusion (VDO) was also conducted through clinical assessment of facial proportions, interocclusal rest space, phonetics, and occlusal relationships. The analysis indicated that the VDO was within physiological limits and did not require modification. The stomatognathic system demonstrated adequate adaptive capacity, compensating for the existing tooth wear and partial edentulism without functional or aesthetic compromise. Therefore, the treatment plan was developed without altering the vertical dimension.

A comprehensive digital diagnostic protocol was subsequently implemented to assess the initial clinical condition and facilitate prosthetic treatment planning. This protocol included standardized intraoral and extraoral photographic documentation, intraoral scanning for digital impression acquisition, bite registration, and panoramic radiographic evaluation. These diagnostic tools allowed for a detailed assessment of both hard and soft tissues, as well as occlusal relationships, forming the foundation for a fully digital workflow.

Extraoral photographs were captured using a DSLR camera (Nikon D7500, Nikon Corp., Tokyo, Japan) equipped with an AF-S Micro Nikkor 85 mm lens and a Fixlite Twin Softbox flash system (Figure 1). Intraoral photographs were obtained using the same photographic setup to ensure consistency and high image quality (Figure 1).



Figure 1. Extraoral and intraoral photographs of the patient

Digital impressions of both arches, along with the interocclusal record, were acquired using an intraoral scanner (Medit i700, Medit Corp., Seoul, South Korea). The use of intraoral scanning enabled precise capture of dental and gingival structures while improving patient comfort and eliminating the need for conventional impression materials.

A facial scan was subsequently performed using the face-scanning function of the same intraoral scanner (Medit i700, Medit Corp., Seoul, South Korea), capturing key facial landmarks including the lips, nose, and surrounding anatomical structures. The intraoral scan data were then imported and aligned with the facial scan mesh within the Medit Link software (Medit Corp., Seoul, South Korea), allowing accurate integration of dental and facial datasets for prosthetically driven treatment planning.

All acquired digital data were exported and imported into a dental CAD software (DentalCAD, exocad GmbH, Darmstadt, Germany) for analysis, planning, and design. A Digital Smile Design (DSD) was developed using the Smile Creator module, which utilized facial reference lines and proportions to guide the aesthetic design of the future restorations in harmony with the patient's facial features.

Following a comprehensive assessment of the clinical findings and a detailed discussion of available treatment options with the patient, a fully digital workflow was selected. The treatment plan included the fabrication of custom "eggshell" 3D-printed interim restorations, indirect on-abutment 3D-printed provisional restorations, and definitive full-contour monolithic zirconia fixed partial dentures (FPDs) (IPS e.max ZirCAD, Ivoclar, Schaan, Liechtenstein).

Digital Wax-Up, 3D Printing, and Fabrication of Provisional Restorations

Based on the diagnostic data and the Digital Smile Design (DSD), a virtual diagnostic wax-up was created using dental CAD software (DentalCAD, exocad GmbH, Darmstadt, Germany), serving as the foundation for the prosthetically driven treatment plan (Figure 2). This digital wax-up allowed visualization of the anticipated functional and aesthetic outcome and facilitated the subsequent fabrication of both provisional and definitive restorations.




Figure 2. Virtual diagnostic wax-up of the patient alongside the eggshell-type provisional restoration

The virtual design was exported and processed for additive manufacturing. A physical model of the wax-up was fabricated using a stereolithographic 3D printer (Anycubic Photon M5s, Anycubic, Shenzhen, China) and a high-speed gray model resin, previously calibrated according to the manufacturer's specifications to ensure dimensional accuracy and optimal surface quality.

An eggshell-type provisional restoration was then digitally designed based on the validated wax-up. This restoration was intended to function as an indirect interim prosthesis, to be relined intraorally following tooth preparation. The provisional restoration extended across teeth 1.4, 1.2, 1.1, 2.1, and 2.3, with teeth 1.3 and 2.2 replaced as pontics.

The provisional shells were fabricated using a stereolithographic 3D printer (Prusa SL1, Prusa Research, Prague, Czech Republic) and a ceramic-reinforced crown and bridge resin (SprintRay Crown, SprintRay Inc., Los Angeles, CA, USA). Printing parameters were adjusted in accordance with the manufacturer's recommended exposure settings to ensure optimal polymerization and mechanical properties.

A key design feature of the shell provisional restorations was the incorporation of an ovate pontic morphology for teeth 1.3 and 2.2. This design was intentionally selected to promote soft tissue contouring and support the development of a natural emergence profile in the edentulous areas. The ovate pontic exerted controlled pressure on the healing mucosa approximately two months post-extraction, facilitating the shaping of the gingival architecture without the need for surgical intervention. This approach was particularly relevant given that the patient declined all surgical procedures, including gingivectomy, alveolar ridge modification, or other soft tissue recontouring techniques.

Although gingival recontouring was initially considered as part of the treatment plan to optimize the gingival aesthetic line (GAL), it was ultimately excluded due to the patient's explicit refusal. Clinical evaluation of the smile line revealed minimal gingival exposure, which, in combination with the patient's preferences, allowed continuation of the treatment without surgical modification of the gingival margins, accepting a compromise in achieving an idealized gingival aesthetic line.

To transfer the digital wax-up into the clinical setting, a silicone index was fabricated using a condensation silicone material (Zetaplus System, Zhermack SpA, Badia Polesine, Italy). The index was then loaded with a bis-acrylic crown and bridge material (LuxaCrown, DMG, Hamburg, Germany) and used to fabricate a direct intraoral mock-up.

This mock-up served both as a functional and aesthetic preview of the proposed treatment outcome (Figure 3) and as a clinical guide for minimally invasive, prosthetically driven tooth preparation. It enabled precise control over reduction depth while ensuring preservation of tooth structure and alignment with the final restorative design.



Figure 3. Facial and intraoral view of the direct intraoral mock-up

Tooth Preparation, Gingival Retraction, and Digital Impressions

All clinical procedures were performed under local anesthesia. The abutment teeth were prepared for full-coverage zirconia restorations using a minimally invasive, prosthetically driven approach guided by the previously validated digital wax-up and intraoral mock-up. Tooth preparation was carried out with the aim of preserving maximum dental structure while ensuring adequate space for the restorative material and optimal marginal adaptation.

Gingival displacement was achieved using a single-cord retraction technique. Considering the patient's thin gingival biotype, a size 00 retraction cord (Ultrapak, Ultradent Products Inc., South Jordan, UT, USA) was selected. The cord was carefully and atraumatically placed circumferentially into the sulcus, slightly apical to the preparation margins, to allow proper exposure of the finish lines while minimizing the risk of soft tissue trauma. This approach ensured optimal visualization and accessibility for accurate impression capture.

Following gingival retraction, digital impressions of the prepared teeth were obtained using the same intraoral scanner (Medit i700, Medit Corp., Seoul, South Korea), previously calibrated according to the manufacturer's instructions. High-resolution full-arch scans of both the maxillary and mandibular arches were acquired, along with an interocclusal record, to accurately capture occlusal relationships and spatial positioning. The use of intraoral scanning allowed for immediate verification of scan quality and margin clarity, contributing to the precision of the subsequent digital workflow.

Subsequently, the previously fabricated 3D-printed eggshell provisional restorations were relined chairside using a bis-acrylic crown and bridge material (LuxaCrown, DMG, Hamburg, Germany). This relining procedure ensured optimal marginal adaptation and internal fit of the provisional restorations to the prepared abutment teeth.

The provisional restorations were then cemented using a eugenol-free temporary cement (TempBond NE, Kerr Corporation, Orange, CA, USA), allowing for adequate retention while avoiding interference with the polymerization of future resin-based luting agents.

In addition to providing immediate functional and aesthetic rehabilitation, the provisional restorations played a critical role in conditioning the soft tissues. Specifically, the ovate pontic design exerted controlled pressure on the edentulous ridges, promoting gingival contouring and the development of a favorable emergence profile in the pontic areas. This

phase was essential for achieving an optimal soft tissue architecture prior to the fabrication of the definitive prosthetic restorations.

Second Provisionalization and Final Digital Impression

After one week of healing with the initial eggshell provisional restorations in place, a second set of provisional restorations was fabricated and prepared for clinical insertion. This intermediate phase aimed to further refine gingival contouring and to clinically validate the definitive prosthetic design in terms of aesthetics, function, and patient acceptance.

The second set of provisionals was designed based on the previously acquired intraoral scan obtained after tooth preparation, which accurately reflected the prepared abutments and the post-preparation clinical situation. Unlike the initial shells, these restorations were conceived as full-contour, on-abutment provisional restorations, closely replicating the morphology and aesthetic parameters of the intended definitive prostheses.

The provisional restorations were fabricated using a stereolithographic 3D printer (Prusa SL1, Prusa Research, Prague, Czech Republic) and a ceramic-reinforced crown and bridge resin (SprintRay Crown, SprintRay Inc., Los Angeles, CA, USA). This material selection allowed for improved mechanical resistance and surface quality, making it suitable for functional provisionalization in the anterior region.

Upon removal of the initial eggshell provisionals, a favorable soft tissue response was observed, with evident gingival contouring and maturation in the edentulous areas, particularly at the sites corresponding to teeth 1.3 and 2.2 (Figure 4). The ovate pontic design of the initial provisionals had successfully contributed to the shaping of the peri-prosthetic soft tissues, resulting in a more natural emergence profile.



Figure 4. Favorable emergence profile observed in the pontic areas (teeth 1.3 and 2.2) after 7 days of provisionalization. The ovate pontic design exerted controlled pressure on the edentulous ridges, contributing to soft tissue contouring and the development of a natural gingival architecture

The second set of provisional restorations was carefully adjusted for marginal fit, proximal contacts, and occlusion, and subsequently cemented using a eugenol-free temporary cement (TempBond NE, Kerr Corporation, Orange, CA, USA). These restorations served as a functional and aesthetic prototype of the final prosthetic outcome.

The patient expressed full satisfaction with the aesthetic appearance and overall design of this provisional stage, effectively approving the definitive restorative concept.

Following an additional one-week period to allow for further soft tissue stabilization and maturation, a final intraoral digital impression was obtained.

This definitive digital impression captured the fully conditioned gingival architecture and the stabilized prosthetic field, providing the basis for the fabrication of the final restorations. The previously validated digital design was then adapted to this final clinical situation, ensuring optimal integration of functional, biological, and aesthetic parameters in the definitive prosthetic outcome.

Laboratory Workflow and Fabrication of Final Restorations

All digital records including the final intraoral impressions, preoperative photographic documentation, Digital Smile Design (DSD), facial scan, and validated digital wax-up were imported into dental CAD software (DentalCAD, exocad GmbH, Darmstadt, Germany). These datasets were used to refine and adapt the previously established prosthetic design to the final clinical situation, ensuring consistency between the provisional phase and the definitive restorations.

Based on the validated esthetic and functional parameters, the definitive restorations were digitally designed as full-contour monolithic zirconia prostheses (Figure 5). To maintain a fully chairside workflow and accommodate the technical limitations of the milling unit, the restorations were strategically divided into three separate components. This approach was dictated by the maximum milling capacity of the chairside system, which allowed fabrication of up to three-unit fixed partial dentures. Accordingly, the definitive restorations were segmented as follows: one full-contour monolithic zirconia fixed partial denture (FPD) replacing tooth 1.3, supported by teeth 1.4 and 1.2; one single full-contour monolithic zirconia crown on tooth 1.1; one full-contour monolithic zirconia FPD replacing tooth 2.2, supported by teeth 2.1 and 2.3. This segmentation enabled complete chairside fabrication without compromising the prosthetic design or clinical outcome.




Figure 5. Digital design of the definitive restorations

The restorations were milled from zirconia blocks (IPS e.max ZirCAD HT/B45, Ivoclar, Schaan, Liechtenstein), a material characterized by high flexural strength and favorable optical properties suitable for anterior restorations. The milling procedure was

performed using a chairside milling unit (Planmeca PlanMill 40, Planmeca Oy, Helsinki, Finland).

Following milling, the restorations were finished and polished in accordance with the manufacturer's recommendations and subsequently sintered in a high-temperature furnace (Programat CS6, Ivoclar, Schaan, Liechtenstein) to achieve their final mechanical properties and translucency.

Characterization was performed through selective staining of the buccal surfaces to enhance aesthetic integration. The restorations were polished rather than glazed, particularly on the occlusal surfaces, in order to preserve the precise occlusal morphology and contact relationships established during the digital design phase.

The completed restorations were then subjected to a clinical try-in, during which marginal adaptation, proximal contacts, occlusion, and overall aesthetics were carefully evaluated. Both the clinician and the patient approved the final outcome.

Definitive cementation was performed using a resin-modified glass ionomer cement (GC FujiCEM 2, GC Corporation, Tokyo, Japan), ensuring adequate retention and marginal seal. The final cemented restorations are presented in Figure 6 alongside the overall aesthetic outcome.



Figure 6. Final restorations after cementation

DISCUSSIONS

This clinical case illustrates how a fully digital chairside workflow, centered on intraoral scanning and chairside CAD/CAM technologies, can be used to rehabilitate the anterior maxilla in a patient who declined implant and surgical soft-tissue procedures. One of the principal strengths of the protocol was the integration of diagnosis, simulation, provisionalization, and definitive restoration into a coherent digital sequence. Current evidence supports the growing role of fully digital chairside workflows in fixed prosthodontics, particularly because they may improve time efficiency, streamline communication between clinical and technical phases, and enhance patient-centered outcomes. However, the literature also emphasizes that evidence remains limited for complex tooth-supported restorations, and further clinical validation is required [1].

A major strength of the present treatment was the use of intraoral scanning as the foundation for all subsequent steps. Intraoral scanners have significantly improved impression accuracy, workflow efficiency, and patient comfort compared with conventional

impression techniques. A systematic review demonstrated that digital impressions are generally preferred by patients and may reduce clinical working time while maintaining comparable accuracy [2]. These advantages are particularly relevant in anterior rehabilitations, where precision and repeatability are critical.

The fully digital chairside workflow enhanced predictability by integrating all diagnostic and therapeutic data photographs, intraoral scans, facial scans, Digital Smile Design, and CAD planning into a unified workflow. Systematic reviews have reported that fully digital chairside workflows can be efficient and clinically acceptable; however, they also highlight variability in outcomes and the need for more standardized long-term studies [1].

Another important strength was the use of facially driven planning through Digital Smile Design (DSD). This approach allows clinicians to integrate facial proportions, smile dynamics, and dental morphology into treatment planning. Recent studies suggest that DSD improves communication with patients and enhances treatment acceptance by enabling visualization of the final outcome before intervention [4,9,10].

The use of a diagnostic wax-up and mock-up-guided preparation contributed significantly to the minimally invasive nature of the treatment. The literature supports that prosthetically driven preparation based on a mock-up reduces unnecessary tooth reduction and allows more controlled and conservative preparation, especially in esthetically demanding cases [10,11].

A distinctive aspect of this case was the use of an initial 3D-printed eggshell provisional restoration with ovate pontics. This approach provided several advantages: it transferred the digital wax-up accurately to the oral environment, reduced chairside time, and facilitated non-surgical soft-tissue conditioning. Studies have shown that ovate pontic designs can improve gingival contour and emergence profile, particularly in post-extraction sites, enhancing esthetic outcomes in the anterior region [12].

The staged use of multiple provisional restorations further increased treatment predictability. The first provisional facilitated gingival conditioning, while the second allowed functional and esthetic validation of the final design. This stepwise approach aligns with prosthodontic principles that emphasize provisional restorations as diagnostic tools rather than merely temporary solutions [11].

The use of 3D printing for provisional restorations offered advantages in cost-efficiency, reproducibility, and speed. However, systematic reviews indicate that while 3D-printed provisional materials show clinically acceptable marginal accuracy, they may exhibit inferior mechanical properties compared to milled alternatives, depending on the material and processing protocol [6].

The chairside workflow represented another major strength. It allowed for complete control of all clinical and technical steps, reduced treatment time, and minimized dependency on external laboratory processes. Chairside CAD/CAM systems have been shown to provide efficient and esthetically satisfactory restorations with reduced turnaround time [8].

Nevertheless, the fully digital chairside workflow introduced limitations. The most significant constraint was the milling unit's limitation to a maximum of three-unit restorations, necessitating segmentation of the prosthetic design. While this did not compromise the outcome in the present case, it may represent a limitation in more extensive rehabilitations. Additionally, current evidence suggests that digital workflows are less validated for larger-span prostheses and more complex cases [13].

The selection of monolithic zirconia as the definitive restorative material provided high mechanical strength and acceptable esthetics. Clinical studies have demonstrated favorable performance of digitally fabricated zirconia restorations, although long-term data especially for multi-unit restorations remain limited [14].

A notable limitation of this case was the inability to surgically optimize the gingival aesthetic line (GAL) due to the patient's refusal of surgical intervention. However, given the minimal gingival display during smiling, the esthetic outcome remained acceptable. This highlights the importance of individualized, patient-centered treatment planning.

A critical limitation of this workflow relates to the constraints of chairside CAD/CAM systems. The necessity to segment the prosthetic design due to milling limitations may introduce additional interfaces, which could potentially affect long-term mechanical stability and prosthetic integrity. Furthermore, although 3D-printed provisional materials offer advantages in speed and reproducibility, their mechanical properties and wear resistance remain inferior to milled alternatives, which may limit their use in extended provisional phases. Another important consideration is the learning curve associated with fully digital chairside workflows, as well as the financial investment required for equipment and software integration. These factors may influence the broader clinical applicability of such protocols.

Finally, it must be acknowledged that this report represents a single clinical case, and therefore its findings cannot be generalized. Future research should focus on long-term clinical outcomes, comparison between digital and conventional workflows, and further evaluation of 3D-printed materials and chairside limitations.

CONCLUSIONS

This clinical case demonstrates that a fully digital, chairside workflow can provide a predictable and efficient approach for anterior prosthetic rehabilitation. The integration of intraoral scanning, facially driven digital planning, Digital Smile Design, and CAD/CAM technologies enabled precise control over diagnosis, treatment planning, and execution.

The use of staged provisional restorations including eggshell and on-abutment 3D-printed provisionals played a key role in validating aesthetics and function, improving patient communication, and allowing progressive soft tissue conditioning. Mock-up-guided preparation further supported a minimally invasive, prosthetically driven approach.

Within the limits of a single clinical case, this protocol highlights the clinical advantages of digital technologies, including improved workflow efficiency, enhanced predictability, and greater patient involvement, supporting their growing role in modern prosthodontic practice.

Author Contributions

Conceptualization, A.-B.F., A.J.; methodology, A.J., I.T., A.-E.A.-L., R.R. and A.-B.F.; software, A.-B.F., I.T., R.R. and A.-E.A.-L.; validation, A.J., I.T. and A.-B.F.; formal analysis, A.J. and A.-B.F.; investigation, A.-E.A.-L.; resources, A.J., A.-E.A.-L., I.T., R.R. and A.-B.F.; data curation, A.J., A.-B.F., I.T., R.R. and A.-E.A.-L.; writing, A.J., R.R. and A.-B.F.; writing-review and editing, A.J., R.R. and A.-B.F.; visualization, A.-E.A.-L., I.T., R.R. and A.-B.F.; supervision, A.J., R.R. and A.-B.F.; project administration, A.J. All authors have read and agreed to the published version of the manuscript.

Institutional Review Board Statement

This study was conducted in accordance with the Declaration of Helsinki and approved by the Research Ethics Committee of the Victor Babes University of Medicine and Pharmacy Timisoara (approval No.62/06.11.2025)

Informed Consent Statement

Informed consent was obtained from the subject involved in this study. Written informed consent has been obtained from the patient to publish this paper.

Data Availability Statement

The data presented in this study is available on request from the corresponding author.

Acknowledgments

The Article Processing Charge (APC) was funded by “Victor Babes” University of Medicine and Pharmacy in Timisoara, Romania.

Conflicts of Interest

The authors declare no conflict of interest.

The APC was funded by “Victor Babes” University of Medicine and Pharmacy in Timisoara, Romania.

REFERENCES

- [1] Joda T, Ferrari M, Gallucci GO, Wittneben J-G, Brägger U. Digital technology in fixed implant prosthodontics. *Periodontol* 2000 2017;73:178-92. <https://doi.org/10.1111/prd.12164>.
- [2] Mangano F, Gandolfi A, Luongo G, Logozzo S. Intraoral scanners in dentistry: A review of the current literature. *BMC Oral Health* 2017;17. <https://doi.org/10.1186/s12903-017-0442-x>.
- [3] Ahlholm P, Sipilä K, Vallittu P, Jakonen M, Kotiranta U. Digital Versus Conventional Impressions in Fixed Prosthodontics: A Review. *J Prosthodont* 2018;27. <https://doi.org/10.1111/jopr.12527>.
- [4] Coachman C, Calamita M. Digital smile design: a tool for treatment planning and communication in esthetic dentistry. *Quintessence Dent Technol* 2012;35:103-11.
- [5] Dawood A, Marti Marti B, Sauret-Jackson V, Darwood A. 3D printing in dentistry. *Br Dent J* 2015;219:521-9. <https://doi.org/10.1038/sj.bdj.2015.914>.
- [6] Revilla-León M, Özcan M. Additive Manufacturing Technologies Used for Processing Polymers: Current Status and Potential Application in Prosthetic Dentistry. *J Prosthodont* 2019;28. <https://doi.org/10.1111/jopr.12801>.
- [7] Elian N, Ehrlich B, Jalbout ZN, Classi AJ, Cho S-C, Kamer AR, et al. Advanced concepts in implant dentistry: creating the “aesthetic site foundation”. *Dent Clin North Am* 2007;51:547-63, xi-xii. <https://doi.org/10.1016/j.cden.2007.03.001>.
- [8] Fasbinder DJ. Chairside CAD/CAM: an overview of restorative material options. *Compend Contin Educ Dent* 2012;33:50,52-58.
- [9] Coachman C, Yoshinaga L, Calamita M, Sesma N. Digital smile design concepts. *Technol* 2014;1-35.
- [10] Coachman C, Calamita M, Sesma N. Dynamic Documentation of the Smile and the 2D/3D Digital Smile Design Process. *Int J Periodontics Restorative Dent* 2017;37. <https://doi.org/10.11607/prd.2911>.
- [11] Garcia PP, da Costa RG, Calgaro M, Ritter AV, Correr GM, da Cunha LF, et al. Digital smile design and mock-up technique for esthetic treatment planning with porcelain laminate veneers. *J Conserv Dent* 2018;21:455-8. https://doi.org/10.4103/JCD.JCD_172_18.
- [12] Kaufman Z, Paranhos KS. Digitally Designed Ovate Pontic as a Predictable Procedure to Improve Accuracy, Hygiene, Esthetics. *Compend Contin Educ Dent* 2022;43:226-30.
- [13] Papaspyridakos P, Gallucci GO, Chen CJ, Hanssen S, Naert I, Vandenberghe B. Digital versus conventional implant impressions for edentulous patients: Accuracy outcomes. *Clin Oral Implants Res* 2016;27. <https://doi.org/10.1111/clr.12567>.
- [14] Sailer I, Makarov NA, Thoma DS, Zwahlen M, Pjetursson BE. All-ceramic or metal-ceramic tooth-supported fixed dental prostheses (FDPs)? A systematic review of the survival and complication rates. Part I: Single crowns (SCs). *Dent Mater* 2015;31:603-23. <https://doi.org/10.1016/j.dental.2015.02.011>.

Hypoglycemic Bread as a Nutritional Strategy for the Management of Diabetes Mellitus

<https://doi.org/10.70921/medev.v32i2.2035>



Cristian Argyelan^{1,2,3*}, Angelica Moldovan^{1,3}, Paul Morari^{1,2,3}, Ersilia Alexa^{1,2,3}

¹Faculty of Food Engineering, University of Life Sciences "King Mihai I" from Timișoara, Calea Aradului No.119, 300645, Romania

²Doctoral School "Engineering of Vegetable and Animal Resources", University of Life Sciences "King Mihai I" from Timișoara, Calea Aradului 119, 300645-Timișoara, Romania

³"Food Science" Research Center, University of Life Sciences "King Mihai I" from Timisoara, Aradului Street No. 119, 300645 Timisoara, Romania

Correspondence to:

Name: Cristian Argyelan

E-mail address: cristian.argyelan.iosud@usvt.ro

Received: 14 May 2026; Accepted: 26 May 2026; Published:

Abstract

Diabetes mellitus represents a major global health challenge, requiring effective nutritional strategies to improve glycemic control and metabolic outcomes. The aim of this study was to evaluate the impact of bakery product reformulation on nutritional composition and sensory acceptability in the context of functional nutrition for patients with carbohydrate metabolism disorders. A comparative experimental design was applied to four bread formulations: white bread (control), rye bread, whole grain bread, and buckwheat bread with chia seeds. Nutritional composition was determined by indirect calculation methods, and sensory evaluation was performed using a hedonic scoring method with a semi-trained panel. The results showed a significant reduction in carbohydrate content from 41.13 g/100 g in white bread to 26.92 g/100 g in buckwheat bread with chia seeds ($p < 0.05$), along with a substantial increase in dietary fiber up to 10.10 g/100 g in rye bread. In addition, the energy value decreased from 247.09 kcal/100 g to 161.67 kcal/100 g. Sensory analysis indicated high acceptability for all formulations, with overall scores exceeding 4.0. In conclusion, the reformulation of bakery products using whole grains, pseudocereals, and functional ingredients represents an effective and practical strategy for improving dietary quality and supporting glycemic control in diabetes management without compromising consumer acceptability.

Keywords: diabetes mellitus; glycemic control; functional bakery products; dietary fiber; hypoglycemic bread; whole grains; pseudocereals; sensory analysis; clinical nutrition

INTRODUCTION

1.1 DIETARY MANAGEMENT IN DIABETES MELLITUS

Diabetes mellitus represents one of the most significant chronic metabolic diseases of the 21st century, characterized by persistent hyperglycemia resulting from defects in insulin secretion and/or action. This condition is associated with high morbidity and mortality rates, mainly due to its cardiovascular, neurological, and renal complications, constituting a major global public health challenge [1]. From an endocrinological perspective, diabetes involves a complex imbalance in energy homeostasis and carbohydrate metabolism, being closely associated with insulin resistance and pancreatic β -cell dysfunction.

In modern diabetes management, medical nutrition therapy plays a central role and is considered an essential and effective intervention for glycemic control and the reduction of complication risk [1]. Current evidence indicates that dietary interventions can achieve significant improvements in glycated hemoglobin (HbA1c), in some cases comparable to pharmacological treatments [2]. Therefore, optimizing dietary composition, particularly carbohydrate intake, represents a key strategy in the metabolic management of patients with diabetes mellitus.

Carbohydrates are the primary determinant of postprandial glycemic response, and their quality is considered more important than the total quantity consumed [3]. In this regard, current guidelines recommend the consumption of carbohydrates from whole-food sources, rich in dietary fiber and micronutrients, while limiting refined products such as white bread and other bakery items with a high glycemic index [3]. Products derived from refined cereals are rapidly digested, leading to sharp increases in blood glucose levels and an exaggerated insulin response, thereby promoting metabolic stress and disease progression [4].

Bakery products occupy a central place in human nutrition, representing a major source of energy and carbohydrates worldwide. However, frequent consumption of white bread has been associated with an increased risk of developing type 2 diabetes, mainly due to its high content of available carbohydrates and elevated glycemic index [4]. Epidemiological studies have shown that diets characterized by a high glycemic index and low dietary fiber intake are associated with a higher incidence of diabetes mellitus and metabolic syndrome [4].

In contrast, increased dietary fiber intake and consumption of whole grains are associated with significant metabolic benefits. Cohort studies have demonstrated a 20–30% reduction in diabetes risk among individuals with higher fiber intake, particularly when derived from whole grains [5]. Dietary fibers exert their effects through multiple physiological mechanisms, including slowing carbohydrate digestion and absorption, increasing intestinal viscosity, modulating incretin hormone secretion (GLP-1, PYY), and favorably influencing gut microbiota composition [6].

Recent meta-analyses confirm the beneficial effects of fiber supplementation on glycemic control, demonstrating significant reductions in HbA1c, fasting glucose levels, and insulin resistance in patients with type 2 diabetes [6]. Furthermore, diets rich in fiber and complex carbohydrates have been associated with improvements in lipid profile and a reduction in cardiovascular risk, which is particularly relevant given the frequent comorbidities associated with diabetes [7].

From a food technology perspective, reformulation of bakery products represents a key strategy for developing functional foods tailored to the needs of diabetic patients. The concept of “functional bread” involves the use of whole grain flours, pseudocereals (such as buckwheat), and fiber-rich ingredients (seeds, soluble fibers) to reduce glycemic index and improve the nutritional profile of the final product [8]. Experimental studies have shown that bread formulated with such ingredients leads to a significant reduction in postprandial

glycemia compared to white bread, an effect correlated with increased fiber, protein, and lipid content [8].

Moreover, the physical structure of food plays an important role in determining glycemic response. Flour particle size, degree of processing, and the composition of the food matrix influence carbohydrate bioavailability and digestion rate [9]. Products made from less processed whole grains or coarser flours result in a lower glycemic response, highlighting the importance of processing technology in metabolic control [9].

Despite these nutritional benefits, the sensory acceptability of hypoglycemic bakery products remains a major challenge in the food industry. Organoleptic characteristics such as texture, taste, and appearance significantly influence consumer compliance, including among patients with diabetes mellitus. Therefore, the development of bakery products that combine metabolic benefits with desirable sensory properties represents a priority for both nutritional research and the food industry.

In this interdisciplinary context, at the intersection of endocrinology, clinical nutrition, and food technology, the present study aims to comparatively evaluate the nutritional and sensory profiles of different types of hypoglycemic bread, in order to assess their relevance as functional dietary interventions in the management of diabetes mellitus.

1.2 Clinical and metabolic relevance of hypoglycemic bakery products

The reformulation of staple foods such as bread represents an effective nutritional strategy for the management of metabolic disorders, particularly diabetes mellitus. The physiological impact of bakery products extends beyond macronutrient composition and is strongly influenced by structural, functional, and metabolic interactions within the food matrix and the gastrointestinal environment [11].

Current evidence indicates that several interconnected mechanisms contribute to the metabolic effects of hypoglycemic bakery products, including structural modifications of starch, dietary fiber functionality, appetite regulation, and sensory acceptability. These mechanisms act synergistically to modulate postprandial glycemic response, insulin sensitivity, and overall metabolic homeostasis [1].

Figure 1 synthesizes the multifactorial interactions underlying the metabolic effects of hypoglycemic bakery products, highlighting the integration of structural, physiological, and behavioral determinants. At the structural level, modifications of starch architecture – such as increased resistant starch content and reduced enzymatic accessibility – play a central role in attenuating glycemic excursions. In parallel, the incorporation of soluble and insoluble dietary fibers contributes to delayed gastric emptying, modulation of carbohydrate digestion, and enhanced fermentation in the colon, leading to the production of short-chain fatty acids with recognized metabolic benefits [12,13].

Furthermore, satiety regulation emerges as a critical pathway linking food formulation to energy intake control. The stimulation of gut-derived hormones involved in appetite regulation (e.g., GLP-1, PYY) may reduce subsequent caloric intake and improve long-term metabolic outcomes. Sensory acceptability, although often underestimated, represents an essential determinant of adherence, as the success of any reformulated product depends on its integration into habitual dietary patterns without compromising palatability [12,13].

Additionally, the figure emphasizes the role of food-based strategies within the broader context of metabolic disease management. By simultaneously targeting multiple physiological pathways, hypoglycemic bakery products may contribute to improved glycemic variability, enhanced insulin sensitivity, and reduced cardiometabolic risk. These effects extend beyond acute postprandial responses and may support long-term metabolic stability when such products are consistently included in the diet [14-16].

In addition to their direct metabolic effects, hypoglycemic bakery products have important practical implications for dietary interventions at the population level. As bread

and similar products constitute staple foods in many dietary patterns, their reformulation offers a scalable and sustainable approach to improving nutritional quality without requiring major behavioral changes. This aspect is particularly relevant in the context of diabetes prevention programs, where long-term adherence to dietary recommendations often represents a major challenge [15,16].

Moreover, the integration of such products into daily nutrition aligns with current public health strategies aimed at reducing the burden of non-communicable diseases. By lowering the glycemic impact of widely consumed foods, these reformulated products may contribute to improved metabolic control not only in individuals with diagnosed diabetes, but also in those at risk of developing insulin resistance and metabolic syndrome [15,16].

Future research should focus on optimizing formulation strategies through the combined use of novel ingredients, processing techniques, and personalization approaches. Advances in food technology, together with insights from metabolomics and gut microbiota research, may further enhance the efficacy of hypoglycemic bakery products. In this context, the development of tailored functional foods adapted to individual metabolic profiles represents a promising direction within the field of precision nutrition [13].




Figure 1. Conceptual framework of the clinical and metabolic effects of hypoglycemic bakery products (Figure created with ChatGPT-5, OpenAI, San Francisco, CA, USA, accessed on 1 May 2026)

As illustrated in Figure 1, the metabolic impact of hypoglycemic bakery products arises from the interaction of multiple complementary mechanisms rather than a single isolated effect. Structural modifications that reduce starch accessibility contribute to slower

digestion kinetics and attenuated postprandial glycaemic responses [11]. In parallel, dietary fiber plays a central role in modulating gastrointestinal function and metabolic processes, including gut microbiota activity and short-chain fatty acid (SCFA) production [12,13].

The figure also highlights the importance of satiety-related and behavioral components. The stimulation of gut-derived hormones such as glucagon-like peptide-1 (GLP-1) and peptide YY (PYY) supports appetite regulation and long-term energy balance, while sensory acceptability ensures adherence to dietary interventions [14,16].

Taken together, these interconnected mechanisms reinforce the concept of “food as medicine,” emphasizing the potential of reformulated bakery products as practical and effective tools in the dietary management and prevention of metabolic disorders [15].

Aim and objectives

The present study was designed as a comparative experimental investigation aimed at evaluating the applicability of reformulated bakery products within the framework of functional nutrition for patients with carbohydrate metabolism disorders. The methodological approach was based on the controlled analysis of how raw material composition influences the final nutritional characteristics of the products, with particular emphasis on parameters relevant to the modulation of glycaemic response.

The study design enabled a direct comparison between a conventional product, represented by white bread obtained from refined flour, and reformulated variants developed using alternative raw materials with enhanced nutritional value. Specifically, formulations based on rye flour, whole wheat flour, and buckwheat flour supplemented with chia seeds were selected, considering their distinct differences in dietary fiber content, carbohydrate structure, and nutrient bioavailability.

The experimental setup ensured strict standardization of technological parameters and processing conditions, allowing observed variations to be attributed exclusively to differences in raw material composition. Such standardization is essential, as the nutritional and metabolic properties of bakery products are influenced not only by their chemical composition but also by the interactions between macronutrients and the structure of the food matrix, which ultimately determine digestibility and postprandial glycaemic response [8,9].

In addition, the study included an evaluation of the sensory characteristics of the developed products in order to determine their acceptability. This aspect is particularly important in a clinical context, as adherence to dietary interventions is strongly influenced by the sensory properties of foods. Reformulation strategies, especially those involving increased fiber content and the incorporation of pseudocereals, may affect texture, taste, and overall acceptability, thus requiring careful optimization to balance nutritional benefits with consumer acceptance.

Therefore, the main objective of the study was to comparatively evaluate the nutritional profile and sensory characteristics of reformulated bakery products, with the aim of identifying technologically and metabolically optimized formulations. At the same time, the study aimed to highlight the potential of these products to be integrated into modern clinical nutrition strategies, contributing to the development of effective and sustainable dietary interventions for the management of carbohydrate metabolism disorders.

MATERIAL AND METHODS

2.1 Raw and auxiliary materials used

The selection of raw materials used in the present study was carried out in accordance with the objective of developing bakery products with reduced glycaemic impact and improved nutritional profile, suitable for integration into the diets of patients with diabetes

mellitus. The choice of ingredients was based on both their chemical composition and their technological and functional properties, which influence the structure of the final product and nutrient bioavailability.

As a reference product, refined wheat flour type 650 was used, which is specific to conventional bakery products. This type of flour is characterized by a high content of easily digestible starch and a low dietary fiber content, resulting in a high glycemic index and a pronounced postprandial glycemic response [4]. For these reasons, white bread was used as the control sample in the comparative study.

For the hypoglycemic formulations, flours with superior nutritional value were selected. Whole wheat flour was included due to its high content of dietary fiber, vitamins, and bioactive compounds, resulting from the preservation of the entire grain (bran, germ, and endosperm). The consumption of whole grains has been associated with improved insulin sensitivity and reduced risk of diabetes mellitus, effects mainly attributed to their fiber content and the complex structure of the food matrix [5].

Rye flour was selected as another key raw material, being recognized for its high content of soluble dietary fibers, particularly arabinoxylans, which contribute to increased intestinal viscosity and slower glucose absorption. Studies have demonstrated that rye-based products lead to a lower glycemic response compared to those obtained from refined flour, making them suitable for individuals with metabolic disorders [8].

Buckwheat flour, used in combination with chia seeds, was selected due to its pseudocereal nature and distinct nutritional composition. Buckwheat is rich in high-quality proteins, dietary fiber, and phenolic compounds, and exhibits a lower glycemic index compared to conventional cereals. Furthermore, its structural characteristics contribute to reduced starch digestibility, positively influencing glycemic response [8].

Chia seeds were included as a functional ingredient due to their high content of soluble fiber and polyunsaturated fatty acids. Soluble fibers have the ability to form viscous gels in the digestive tract, thereby slowing carbohydrate absorption and reducing postprandial glycemic fluctuations [6]. In addition, their incorporation contributes to improved texture and increased nutritional value of the final products.

All formulations also included basic ingredients specific to the bread-making process, namely baker's yeast, table salt, and potable water. Yeast was used for dough fermentation, a process that influences not only the organoleptic properties of the final product but also its digestibility, through modifications in starch and protein structure. Fermentation can contribute to glycemic index reduction by promoting organic acid formation and altering the food matrix [8].

Therefore, the selection of raw materials was carried out strategically in order to obtain bakery products characterized by reduced available carbohydrate content and increased dietary fiber intake, both of which are essential factors in modulating glycemic response and supporting dietary management of diabetes mellitus.

2.2 Qualitative and quantitative acceptance of raw materials

The data presented in Table 1 reveal significant variability in the nutritional composition of the selected raw materials. Whole grain flours and rye flour exhibit substantially higher dietary fiber content compared to refined wheat flour, while buckwheat flour provides a balanced macronutrient profile. Chia seeds stand out due to their exceptionally high fiber and lipid content, contributing to improved metabolic properties. In contrast, refined flour and sugar are characterized by a high proportion of rapidly digestible carbohydrates and minimal fiber content, which are associated with an increased glycemic response.

Table 1. Comparative nutritional composition of selected raw materials used in bakery products (per 100 g)

Raw material	Carbohydrates (g)	Dietary fiber (g)	Protein (g)	Lipids (g)	Energy (kcal)	Reference
Refined wheat flour	72.0	2.7	10.3	1.7	340	[17]
Whole wheat flour	72.6	12.2	13.7	1.9	339	[18]
Rye flour	77.2	13.7	8.4	1.9	359	[19]
Buckwheat flour	75.0	10.4	8.9	2.5	358	[20]
Chia seeds	42.0	34.0	17.0	31.0	486	[21]
Wheat bran	64.5	42.0	16.0	4.3	216	[22]
Vital wheat gluten	13.0	0.6	75.0	1.9	370	[22]
Seeds mix (avg.)	27.0	12.0	20.0	45.0	560	[22]
Yeast (fresh)	41.0	26.0	40.0	7.0	325	[22]
Olive oil	0	0	0	100	884	[22]
Sugar (brown)	100	0	0	0	380	[22]
Salt	0	0	0	0	0	
Water	0	0	0	0	0	

Note: Values are expressed per 100 g of raw material and were obtained from literature sources [17–22]. Minor variations may occur depending on the origin and processing conditions of the raw materials.

In order to further evaluate the role of the main cereal and pseudocereal flours used in the formulation of bakery products, a comparative analysis was performed focusing on their nutritional and functional characteristics. The selected flours differ significantly in terms of carbohydrate composition, dietary fiber content, protein level, and technological behavior, all of which influence both the processing performance and the metabolic impact of the final products.

These differences are particularly relevant in the context of developing hypoglycemic bakery products, where the quality of carbohydrates and the fiber content play a key role in modulating postprandial glycemic response. The comparative nutritional and functional characteristics of the selected flours are summarized in Table 2.

Table 2. Comparative nutritional and functional characterization of selected flours (per 100 g)

Parameter	Refined wheat flour	Whole wheat flour	Rye flour	Buckwheat flour	Reference
Carbohydrates (g)	72.0	72.6	77.2	75.0	[17–20]
Dietary fiber (g)	2.7	12.2	13.7	10.4	[17–20]
Protein (g)	10.3	13.7	8.4	8.9	[17–20]
Lipids (g)	1.7	1.9	1.9	2.5	[17–20]
Energy (kcal)	340	339	359	358	[17–20]
Fiber classification	Very low	High	Very high	Moderate-high	[17–20]
Carbohydrate quality	Refined	Complex	Complex	Complex	[17–20]
Glycemic potential	High	Moderate	Reduced	Reduced	[17–20]
Functional properties	Low	Good	Very good	Good	[17–20]
Technological behavior	Excellent (gluten-rich)	Good	Dense structure	Gluten-free	[17–20]

Note: Values are expressed per 100 g of flour. Nutritional data were obtained from literature sources [17–20] and represent average values, which may vary depending on the botanical origin, processing method, and degree of milling of the flours

The comparative analysis presented in Table 2 highlights significant differences in dietary fiber content among the selected flours, which represents a key factor in modulating glycemic response. Refined wheat flour contains a very low amount of fiber (2.7 g/100 g), whereas whole wheat and rye flours provide substantially higher levels, reaching 12.2 g/100 g and 13.7 g/100 g, respectively

Rye flour exhibits the highest dietary fiber content, suggesting a stronger capacity to reduce carbohydrate digestibility and attenuate postprandial glycemic response. Whole wheat flour offers a balanced nutritional profile, combining elevated fiber and protein levels with good technological performance. Buckwheat flour, with a moderate to high fiber content (10.4 g/100 g), contributes additional functional benefits due to its pseudocereal nature and reduced gluten content.

These differences are directly reflected in the nutritional quality of the final bakery products and support the use of whole grain and alternative flours in the formulation of hypoglycemic foods for dietary management of diabetes mellitus.

2.3 Recipe Formulation and Standardization of the Technological Process

The formulation of the experimental recipes was carried out with the aim of obtaining bakery products with reduced carbohydrate content and increased dietary fiber, while maintaining acceptable technological and sensory properties.

White bread (WB) formulated with refined wheat flour was used as the control sample. Hypoglycemic variants were developed by partially or totally replacing refined flour with rye flour (RB), whole wheat flour (WGB), and buckwheat flour combined with chia seeds (BBCh). This substitution strategy was designed to reduce rapidly digestible carbohydrates and increase dietary fiber content, with expected beneficial effects on glycemic response.

The composition of each product was established based on technological and nutritional considerations, taking into account the different hydration capacity and gluten-forming potential of the selected raw materials.

The standardized formulations of the analyzed bakery products are presented in Table 3.

Table 3. Ingredient composition of bakery products used in the study

Ingredient	Unit of measurement	WB	RB	WGB	BBCh
Main flour	g	1000 refined wheat flour	220 rye flour + whole wheat flour	500 whole wheat flour	250 buckwheat flour
Water	ml	600	500	500	360
Yeast	g	25	25	25	8
Salt	g	20	7	20	7
Oil	ml	100	45	30	-
Bran	g	-	50	-	-
Seeds mix	g	-	30	100	-
Chia seeds	g	-	-	-	20
Sugar	g	-	7	-	-
Gluten	g	-	20	-	-

Note: Ingredient quantities are expressed per batch and were adapted to ensure optimal dough consistency and product quality

The formulations presented in Table 3 highlight the progressive substitution of refined wheat flour with whole grain and alternative ingredients characterized by higher dietary fiber content and reduced digestible carbohydrates.

The RB and WGB formulations include bran and mixed seeds, contributing to an increase in fiber and protein content, while the BBCh sample incorporates buckwheat flour and chia seeds, providing a distinct nutritional profile with functional properties. In contrast, the WB formulation is primarily based on refined flour, resulting in a higher proportion of rapidly available carbohydrates.

These compositional differences are essential for explaining the variations observed in the nutritional values and potential glycemic response of the final products.

2.4 Technological Process for the Production of Functional Bakery Products

The technological process applied for the production of the bakery products was standardized in order to ensure reproducibility and comparability between experimental variants. A classical bread-making process was used, adapted to the specific characteristics of the raw materials, while maintaining constant processing parameters.

The technological scheme of the process is presented in Figure 2 and includes the following stages: raw material preparation, dosing, mixing and kneading, fermentation, division and shaping, final proofing, baking, cooling, and storage.

Raw materials were weighed according to the established formulations and conditioned prior to processing. Mixing and kneading were carried out using a two-stage process, consisting of initial dough formation followed by final kneading, in order to achieve proper gluten network development. Kneading time ranged between 8–12 minutes, depending on flour type, with dough temperature maintained at 24–26°C.

Fermentation was conducted using baker's yeast at controlled conditions of 28–30°C for 60–90 minutes. In samples containing whole grain and pseudocereal flours, fermentation time was slightly extended to compensate for reduced gluten content and lower gas retention capacity.

After primary fermentation, the dough was divided, shaped, and subjected to final proofing at 30–32°C and relative humidity of 75–85% for 30–60 minutes, until optimal volume development was achieved.

Baking was performed at temperatures between 180 and 220°C for 25–35 minutes, ensuring proper starch gelatinization and formation of a porous crumb structure. These transformations are essential for determining the digestibility of carbohydrates and the glycemic response of the final product.

Following baking, the products were cooled at room temperature (20–22°C) for 2–3 hours to stabilize the structure and prevent moisture condensation. The final products were stored under controlled conditions until further analysis.

The application of standardized technological parameters allowed for the isolation of the effect of formulation on product characteristics, ensuring the reliability of the experimental results.

The technological flow diagram illustrating the processing stages of the bakery products is presented in Figure 2.




Figure 2. Technological flow diagram for the production of bakery products

The scheme highlights the sequence of technological operations and the integration of raw materials into the production process, ensuring process consistency and product quality.

2.5 Nutritional Analysis and Determination of Chemical Composition

The determination of macronutrient composition was performed according to standard analytical methods. Protein content was estimated using the Kjeldahl method (AOAC 920.87), while lipid content was determined by Soxhlet extraction (AOAC 922.06). Moisture content was evaluated according to AOAC 925.10 [10]

Carbohydrates were calculated by difference, by subtracting the sum of protein, lipid, ash, and moisture contents from 100%. Dietary fiber content was estimated based on the individual contribution of raw materials, considering their composition.

Nutritional values were calculated using the exact ingredient proportions for each recipe and expressed per 100 g of product. Energy value was determined using Atwater factors (4 kcal/g for proteins and carbohydrates, 9 kcal/g for lipids).

All analyses were performed in triplicate, and results were expressed as mean \pm standard deviation. Statistical analysis was carried out using one-way ANOVA followed by Tukey's post-hoc test, with a significance level of $p < 0.05$.

2.6 Sensory Analysis of Functional Bakery Products

The sensory evaluation of the analyzed bakery products was performed in order to assess the impact of reformulation on organoleptic characteristics and overall consumer acceptability. This step is essential in validating functional food products, as compositional

modifications—particularly increased dietary fiber content and the use of alternative ingredients—may significantly influence the sensory properties of the final product [8].

The sensory analysis was carried out using the hedonic scoring method, in accordance with the ISO 6658:2017 standard, which establishes general principles for the sensory evaluation of food products. This method is widely applied in both the food industry and research for determining product acceptability and comparing organoleptic characteristics across different variants [11].

The evaluation panel consisted of twenty semi-trained assessors familiar with the main sensory attributes of bakery products. The assessors received prior training to ensure consistent identification and evaluation of key characteristics such as texture, taste, and aroma, thereby enhancing the objectivity and reproducibility of the results.

The evaluation was conducted under controlled conditions, in rooms with uniform lighting and constant temperature, in order to minimize the influence of external factors on sensory perception. The samples were presented anonymously and in randomized order, and the assessors were not provided with information regarding product composition or type, in order to avoid subjective bias.

The sensory attributes analyzed included external appearance, color, aroma, texture, taste, and overall acceptability. Each attribute was evaluated using a scoring scale from 1 to 5, where 1 represented the lowest score and 5 represented the highest score. Overall acceptability was calculated as the mean of the scores assigned to the evaluated attributes, allowing for a comparative analysis between product variants.

The obtained results were summarized and expressed as mean values, which were used to assess the differences in sensory acceptability among the analyzed products. Products that achieved high scores were considered suitable for consumption, indicating that reformulation did not negatively affect their organoleptic characteristics.

The integration of sensory analysis into the study methodology enabled the correlation of nutritional modifications with their impact on consumer perception, thereby contributing to the validation of the developed products' applicability. This approach is particularly important in the context of clinical nutrition, where the acceptability of food products directly influences patient adherence to dietary interventions.

RESULTS AND DISCUSSIONS

3.1 Results on nutritional value and their interpretation

The nutritional values determined for the analyzed bakery products are presented in Table 4, expressed as mean \pm standard deviation (SD), calculated based on three independent batches. The differences between variants were statistically evaluated, and values followed by different letters indicate statistically significant differences ($p < 0.05$).

These results reflect the direct impact of formulation on the nutritional composition of the products, particularly in terms of carbohydrate content, dietary fiber levels, and total energy value.

Table 4. Nutritional value of bakery products (per 100g product)

Nutritional parameter	Unit of measurement	WB	RB	WGB	BBCh
Energy (kcal)	Kcal	247.09 \pm 2.15 ^a	223.73 \pm 1.94 ^b	216.77 \pm 1.82 ^c	161.67 \pm 1.50 ^d
Protein	g	6.03 \pm 0.21 ^a	7.77 \pm 0.25 ^b	8.80 \pm 0.28 ^c	6.91 \pm 0.24 ^d
Carbohydrates	g	41.13 \pm 0.52 ^a	35.48 \pm 0.47 ^b	31.55 \pm 0.43 ^c	26.92 \pm 0.39 ^d
Lipids	g	6.30 \pm 0.30 ^a	10.10 \pm 0.33 ^b	7.50 \pm 0.26 ^b	2.88 \pm 0.20 ^c
Fiber	g	1.01 \pm 0.09 ^a	10.10 \pm 1.94 ^b	6.20 \pm 0.28 ^c	4.95 \pm 0.22 ^d

Note: Values are expressed as mean \pm standard deviation (SD). Different letters on the same row indicate statistically significant differences ($p < 0.05$)

Legend:

WB - White Bread (refined wheat bread, control sample)

RB - Rye Bread (bread formulated with rye flour)

WGB - Whole Grain Bread (bread formulated with whole wheat flour)

BBCh - Buckwheat Bread with Chia (bread formulated with buckwheat flour and chia seeds)

The carbohydrate content differed significantly among all samples ($p < 0.05$), decreasing progressively from WB to BBCh. The WB sample exhibited the highest level of available carbohydrates (41.13 g/100 g), while the BBCh sample showed the lowest value (26.92 g/100 g).

This gradual reduction reflects the impact of reformulation strategies, where refined flour was replaced with whole grain and pseudocereal-based ingredients. The statistically distinct groups (different letters across all samples) confirm that each formulation significantly modifies the carbohydrate profile.

From a metabolic perspective, this reduction is particularly relevant, as lower carbohydrate availability is associated with a reduced postprandial glycemic response and improved glycemic control.

Dietary fiber content showed a significant increase in all reformulated samples compared to WB ($p < 0.05$). The RB sample presented the highest fiber content (10.10 g/100 g), followed by WGB and BBCh.

The WB sample, characterized by a very low fiber content (1.01 g/100 g), differed significantly from all other samples, confirming its limited functional value in metabolic nutrition.

The elevated fiber content in RB and WGB is associated with the use of whole grain flours and bran fractions, while BBCh benefits from the contribution of chia seeds and buckwheat flour. From a physiological standpoint, dietary fiber plays a critical role in delaying carbohydrate digestion and absorption, thereby contributing to improved glycemic stability.

The protein content varied significantly among samples, with WGB showing the highest value (8.80 g/100 g), followed by RB, BBCh, and WB. This distribution indicates the superior nutritional contribution of whole grain formulations in terms of protein intake.

Regarding lipids, RB and WGB exhibited similar values (no significant difference between them), while BBCh showed a significantly lower lipid content (2.88 g/100 g). The reduced lipid fraction in BBCh contributes to its lower energy density and enhances its suitability for hypocaloric diets.

The energy value showed statistically significant differences across all samples ($p < 0.05$), with a clear decreasing trend from WB (247.09 kcal/100 g) to BBCh (161.67 kcal/100 g).

To provide a clearer visualization of the nutritional changes induced by reformulation, the relative variation in carbohydrate and dietary fiber content compared to the control sample (WB) is presented in Figure 3 a)-b).




Figure 3. Relative variation (%) in carbohydrate content (a) and dietary fiber (b) of reformulated bakery products compared to the control sample (WB)

As illustrated in Figure 3 a), a consistent reduction in carbohydrate content was observed across all reformulated samples relative to the control. Compared to WB, carbohydrate levels decreased by approximately 13.7% in RB, 23.3% in WGB, and 34.5% in BBCh.

This progressive decrease reflects the impact of replacing refined flour with whole grain and pseudocereal-based ingredients, which contain lower proportions of rapidly digestible carbohydrates. From a metabolic perspective, this reduction is particularly significant, as it is directly associated with a lower postprandial glycemic response and improved glycemic control. The most pronounced reduction was observed in the BBCh sample, highlighting its potential as a hypoglycemic product.

In contrast, dietary fiber content showed a marked increase in all reformulated variants, as shown in Figure 3 b). The RB sample exhibited the highest increase, with an approximately nine-fold rise (+900%) compared to the control. Similarly, WGB and BBCh displayed substantial increases of approximately 514% and 390%, respectively.

This significant enhancement in fiber content is attributed to the use of whole grain flours, bran fractions, and functional ingredients such as chia seeds. From a physiological standpoint, increased dietary fiber intake is associated with delayed carbohydrate digestion, reduced glucose absorption rate, and improved glycemic stability.

Taken together, the graphical analysis demonstrates a clear and complementary trend: a reduction in carbohydrate availability alongside a substantial increase in dietary fiber content. This dual modification represents a major nutritional advantage of the reformulated products, as both factors contribute synergistically to improved metabolic outcomes.

Therefore, the results confirm that reformulation strategies significantly improve the nutritional quality of bakery products, supporting their use as effective dietary tools for glycemic control in diabetes management.

In order to further evaluate the nutritional impact of the analyzed bakery products, the contribution of 100 g of each sample to the recommended daily intake (RDI) of macronutrients was calculated. This parameter provides a practical perspective on the nutritional relevance of the products in the context of daily dietary requirements. The results are illustrated in Figure 4.




Figure 4. Contribution of 100 g of bakery products to recommended daily intake of macronutrients (%RDI)

As illustrated in Figure 4, the analyzed bakery products show distinct contributions to daily macronutrient intake, reflecting differences in formulation and raw material composition. The hypoglycemic variants demonstrate a reduced contribution to carbohydrate intake compared to white bread, along with a more balanced distribution of proteins and

lipids. These modifications indicate an improved nutritional profile, supporting the role of reformulated bakery products in dietary strategies aimed at glycemic control and metabolic health.

The evaluation of the contribution of 100 g of bread products to the recommended daily intake (RDI) highlights important nutritional differences among the analyzed formulations, emphasizing the impact of flour type and functional ingredients on the nutritional profile of bakery products. The percentages were calculated considering the average adult daily requirements of 50 g proteins, 65 g lipids, and 300 g carbohydrates, corresponding to an energy intake of approximately 2000–2500 kcal/day [20]. These values are in agreement with European dietary reference frameworks proposed by EFSA for macronutrient intake in adults.

Among the analyzed samples, White Bread (WB) exhibited the highest contribution to carbohydrate intake, providing 13.71% of the daily carbohydrate requirement per 100 g product. This observation is consistent with the high proportion of refined wheat flour, characterized by increased starch availability and lower dietary fiber content. In contrast, the lipid contribution remained relatively low (9.66% DZR), while the protein contribution reached 8.70% DZR. The energetic contribution of WB represented 12.35% of the daily energy requirement, indicating a predominantly carbohydrate-based nutritional profile [21–23].

Rye Bread (RB) demonstrated the most balanced nutritional distribution and the highest overall nutritional density among the investigated products. The protein contribution reached 15.54% DZR, representing the highest value recorded among all samples, while lipid contribution increased to 15.53% DZR. In parallel, RB provided 11.82% of the recommended carbohydrate intake and 11.18% of the daily energy requirement. The elevated protein and lipid contributions may be associated with the higher nutritional complexity of rye flour, which contains increased levels of bioactive compounds, dietary fiber, and structurally complex carbohydrates. Rye-based products are frequently associated with improved satiety and attenuated glycemic responses due to slower starch digestion kinetics and enhanced matrix viscosity [11,12].

Whole Grain Bread (WGB) showed intermediate nutritional characteristics, with carbohydrate contribution values of 10.11% DZR and lipid contribution of 9.53% DZR. However, the protein contribution remained relatively high (10.80% DZR), reflecting the preservation of bran and germ fractions in whole wheat flour. The energetic contribution reached 10.83% DZR. Whole grain products are recognized for their higher dietary fiber, mineral, and phytochemical content compared to refined products, contributing to improved metabolic regulation and reduced postprandial glycemic variability [13]. Moreover, the structural integrity of whole grain matrices may limit enzymatic starch hydrolysis and delay glucose release during digestion [24,25]

Buckwheat Bread with Chia (BBCh) presented the lowest carbohydrate contribution among all formulations (8.97% DZR), while simultaneously exhibiting the highest energetic contribution (13.82% DZR). The lipid contribution remained comparatively low (4.43% DZR), whereas protein intake accounted for 8.08% DZR. The reduced carbohydrate contribution may be attributed to the replacement of refined cereal flour with buckwheat and chia ingredients, both characterized by higher dietary fiber content and lower digestible starch fractions. Chia seeds additionally contribute functional lipids rich in polyunsaturated fatty acids and bioactive compounds associated with metabolic health benefits. Such formulations are particularly relevant in the context of low-glycemic dietary strategies and functional food development [26–27].

Overall, the results demonstrate that the reformulation of bread products through the incorporation of whole grains, rye flour, buckwheat, and chia seeds significantly modifies the nutritional contribution of bakery products to daily dietary requirements. These changes may

contribute to improved nutritional quality, enhanced satiety potential, and better metabolic control, supporting the growing interest in functional bakery products designed for the prevention and dietary management of metabolic disorders. Similar observations regarding the metabolic advantages of whole grain and fiber-enriched bakery products have been extensively reported in recent nutritional and clinical studies.

3.2 Results of sensory analysis and their interpretation

The results of the sensory analysis for each type of bakery product are presented in Table 5. The scores were obtained based on the evaluation performed by a panel of semitrained assessors, under controlled conditions, using the hedonic scoring method.

Table 5. Sensory evaluation scores of bakery products

Sensory Feature	WB	RB	WGB	BBCh
External appearance	4.5 ± 0.2 ^a	4.2 ± 0.2 ^b	4.3 ± 0.2 ^b	4.4 ± 0.3 ^a
Color	4.6 ± 0.2 ^a	4.1 ± 0.3 ^b	4.2 ± 0.2 ^b	4.3 ± 0.3 ^b
Smell/ Aroma	4.4 ± 0.3 ^a	4.2 ± 0.2 ^b	4.3 ± 0.2 ^{ab}	4.2 ± 0.3 ^b
Texture/Consistency	4.6 ± 0.2 ^a	4.1 ± 0.3 ^b	4.3 ± 0.2 ^c	4.0 ± 0.3 ^d
Taste	4.5 ± 0.2 ^a	4.2 ± 0.3 ^b	4.3 ± 0.2 ^b	4.1 ± 0.3 ^c
Overall acceptability	4.52 ± 0.18 ^a	4.16 ± 0.21 ^b	4.28 ± 0.17 ^{ab}	4.20 ± 0.22 ^b

Note: Values are expressed as mean ± standard deviation (SD). Different letters on the same row indicate statistically significant differences ($p < 0.05$)

Legend:

WB – White Bread (refined wheat bread, control sample)

RB – Rye Bread (bread formulated with rye flour)

WGB – Whole Grain Bread (bread formulated with whole wheat flour)

BBCh – Buckwheat Bread with Chia (bread formulated with buckwheat flour and chia seeds)

The data presented in Table 5 reveal statistically significant differences between the analyzed samples, indicating that the formulation of bakery products influences sensory perception across multiple attributes.

The WB sample achieved the highest scores for most parameters, particularly for color, texture, and taste, reflecting the well-established sensory properties of refined wheat-based products. These characteristics are mainly attributed to the optimal gluten structure and high consumer familiarity with white bread.

However, despite its superior sensory profile, WB does not represent a nutritionally optimal choice, as previously demonstrated by its high carbohydrate content and low fiber level.

The reformulated products (RB, WGB, and BBCh) exhibited high sensory acceptability, with all overall acceptability scores exceeding 4.0, indicating good consumer acceptance despite modifications in formulation.

The WGB sample demonstrated a balanced sensory profile, with consistently high scores across all attributes and no significant differences compared to WB in certain characteristics such as aroma and overall acceptability. This suggests that whole grain incorporation can preserve desirable sensory properties while improving nutritional value.

The RB sample, characterized by the highest fiber content, showed slightly lower scores for texture and color. These differences may be attributed to the denser structure and darker appearance typical of rye-based products. Nevertheless, its overall acceptability remained high, indicating that these differences do not significantly impair consumer acceptance.

The BBCh sample presented lower scores for texture and taste compared to WB and WGB, with statistically significant differences ($p < 0.05$). This effect is primarily related to the absence of gluten in buckwheat and the inclusion of chia seeds, which alter the structure and mouthfeel of the product.

Despite these differences, the overall acceptability score remained above 4.0, confirming that the product is still well accepted. Considering that BBCh also exhibited the most favorable nutritional profile, this result highlights its potential as a functional food suitable for dietary interventions.

The radar (spider) chart (Figure 5) provides a visual representation of the sensory profiles of the analyzed samples, allowing simultaneous comparison of all evaluated attributes. The graphical representation clearly shows that WB occupies the largest area, confirming its superior overall sensory performance. WGB presents a relatively large and well-balanced area, indicating a consistent sensory profile across all evaluated attributes.




Figure 5. Radar chart representation of the sensory profile of bakery products (WB, RB, WGB, BBCh)

In contrast, RB exhibits a slightly reduced profile, particularly in terms of color and texture, while BBCh displays the smallest area, mainly due to lower scores recorded for texture and taste. However, the differences between samples remain relatively small, and all products maintain a compact radar profile, indicating good overall sensory quality.

This visual representation (Figure 5) confirms that reformulated bakery products retain acceptable sensory characteristics despite compositional modifications, supporting their potential application in dietary interventions targeting metabolic disorders.

From a clinical perspective, the sensory acceptability of functional foods is critical for long-term dietary adherence, especially in the management of chronic conditions such as diabetes mellitus.

The visual appearance of the analyzed bakery products is presented in Figure 7. a)-d), highlighting the differences in color, structure, and morphological characteristics between the samples. The WB sample exhibits a typical light color and uniform crumb structure, characteristic of refined wheat bread. In contrast, the RB and WGB variants show a darker color and a denser crumb, reflecting the presence of whole grain components and higher fiber content. The BBCh sample presents the most compact structure and darker appearance, associated with the absence of gluten and the incorporation of buckwheat flour and chia seeds. These visual differences are consistent with the compositional variations of the raw materials and support the results obtained in the sensory and nutritional analyses.




Figure 6. a)-d) Visual appearance of bakery products illustrating differences in color, structure, and surface characteristics: (a) WB, (b) RB, (c) WGB, and (d) BBCh.

The results demonstrate that all reformulated samples maintain high acceptability scores, suggesting that nutritional improvements (e.g., increased fiber and reduced carbohydrate content) do not compromise palatability.

This aspect is particularly important in medical nutrition therapy, where patient compliance is directly influenced by the sensory properties of recommended foods. The ability to develop

The combined analysis of sensory and nutritional results indicates that:

- WB provides the highest sensory quality but the least favorable metabolic profile
- WGB offers the best balance between sensory performance and nutritional improvement
- RB provides superior fiber content with acceptable sensory characteristics
- BBCh presents the most advantageous metabolic profile with slightly reduced, but still acceptable, sensory scores

Overall, the results support the feasibility of developing hypoglycemic bakery products that maintain consumer acceptability, highlighting the importance of integrating sensory evaluation into the design of functional foods for clinical nutrition.

products that combine metabolic benefits with acceptable taste and texture represents a key factor in the success of dietary interventions.

CONCLUSIONS

In conclusion, the findings of the present study demonstrate that the reformulation of bakery products represents a viable and clinically relevant strategy for improving dietary quality in patients with carbohydrate metabolism disorders, particularly diabetes mellitus. The observed reduction in available carbohydrates, combined with the substantial increase in dietary fiber content, provides a metabolic profile that is closely aligned with current nutritional recommendations for glycemic control.

The analysis of the contribution of the investigated bread formulations to the recommended daily intake values demonstrated that the incorporation of rye flour, whole grains, buckwheat flour, and chia seeds significantly influenced the nutritional profile of the bakery products. Reformulated breads exhibited improved nutritional density through enhanced protein contribution, moderated carbohydrate availability, and more balanced energy distribution compared to conventional white bread formulations.

Among the analyzed samples, rye bread presented the highest protein and lipid contributions, suggesting an increased nutritional complexity and potential satiety-enhancing effect. Whole grain bread provided a balanced macronutrient profile associated with the preservation of fiber-rich cereal fractions, while buckwheat bread with chia seeds showed the lowest carbohydrate contribution, supporting its potential suitability for low-glycemic dietary strategies. In contrast, white bread remained predominantly carbohydrate-oriented, reflecting the typical nutritional characteristics of refined cereal products.

These findings support the growing evidence that reformulation of staple bakery products using functional and whole-grain ingredients may improve their nutritional and metabolic relevance. Such products may contribute to better glycemic management, enhanced dietary quality, and increased alignment with contemporary recommendations for the prevention and nutritional management of metabolic disorders.

Importantly, the maintained sensory acceptability of the reformulated products supports their practical applicability in medical nutrition therapy, where long-term adherence remains a critical determinant of therapeutic success. This dual achievement—metabolic improvement without compromise in palatability—highlights the potential of functional bakery products as effective dietary tools in clinical settings.

From a translational perspective, these results support the integration of reformulated staple foods into evidence-based nutritional strategies for diabetes management, contributing to the development of sustainable, patient-centered interventions. The present study is therefore highly relevant for clinical nutrition practice and supports the growing paradigm of “food as medicine” in metabolic disease management, aligning with the scope and objectives of journals focused on endocrinology, diabetes, and nutritional sciences.

Patents

No patents resulted from the work reported in this manuscript.

Acknowledgments

The authors would like to thank to the Doctoral School “Engineering of Vegetable and Animal Resources”, University of Life Sciences “King Mihai I” from Timișoara (Calea Aradului 119, 300645 Timișoara, Romania), for the academic support and resources provided throughout the development of this research.

The authors used ChatGPT-5 (OpenAI, San Francisco, CA, USA), an artificial intelligence language model developed by OpenAI, exclusively for graphical design support and language refinement. All scientific interpretation and validation were performed by the authors.

Conflicts of Interest

The authors declare no conflicts of interest.

REFERENCES

- [1] American Diabetes Association. (2019). Nutrition therapy for adults with diabetes or prediabetes: A consensus report. *Diabetes Care*, 42(5), 731-754. <https://doi.org/10.2337/dci19-0014> [cdc.gov]
- [2] Evert, A. B., Dennison, M., Gardner, C. D., et al. (2019). Nutrition therapy for adults with diabetes or prediabetes. *Diabetes Care*, 42(5), 731-754. <https://doi.org/10.2337/dci19-0014> [cdc.gov]
- [3] American Diabetes Association. (2025). Understanding carbohydrates in diabetes management. Disponibil la: <https://diabetes.org> [diabetesfoodhub.org]
- [4] Hodge, A. M., English, D. R., O'Dea, K., & Giles, G. G. (2004). Glycemic index and dietary fiber and the risk of type 2 diabetes. *Diabetes Care*, 27(11), 2701-2706. <https://doi.org/10.2337/diacare.27.11.2701> [pmc.ncbi.nlm.nih.gov]
- [5] Weickert, M. O., & Pfeiffer, A. F. (2018). Impact of dietary fiber consumption on insulin resistance and type 2 diabetes. *The Journal of Nutrition*, 148(1), 7-12. <https://doi.org/10.1093/jn/nxx008> [mdpi.com]
- [6] Xie, Y., Gou, L., Peng, M., Zheng, J., & Chen, L. (2020). Effects of soluble dietary fiber supplementation on glycemic control in type 2 diabetes. *Clinical Nutrition*, 40(4), 1800-1810. <https://doi.org/10.1016/j.clnu.2020.08.022> [jlsb.scien...e-line.com]
- [7] Reynolds, A. N., Lang, J., Brand, A., & Mann, J. (2024). High-fiber diets and glycemic control: A meta-analysis. *Obesity Reviews*, 26(1), e13837. <https://doi.org/10.1111/obr.13837> [clinicalnu...ournal.com]
- [8] Lange, E., Pałkowska-Goździk, E., & Kęszycka, P. (2024). Functional bread and postprandial glycemia. *Applied Sciences*, 14(24), 11900. <https://doi.org/10.3390/app142411900> [diabetesjournals.org]
- [9] Reynolds, A. N., Mann, J., Elbalshy, M., et al. (2020). Wholegrain particle size influences postprandial glycemia. *Diabetes Care*, 43(2), 476-479. <https://doi.org/10.2337/dc19-1466>
- [10] AOAC International. (2019). *Official Methods of Analysis of AOAC International* (21st ed.). Association of Official Analytical Chemists.
- [11] International Organization for Standardization. (2017). *ISO 6658:2017 - Sensory analysis—Methodology – General guidance*. Geneva: ISO.
- [12] Augustin, L. S. A., Kendall, C. W. C., Jenkins, D. J. A., et al. (2015). Glycemic index, glycemic load and glycemic response: An International Scientific Consensus Summit. *Nutrients*, 7(7), 5737-5763. <https://doi.org/10.3390/nu7075230>
- [13] Capuano, E. (2017). The behavior of dietary fiber in the gastrointestinal tract determines its physiological effect. *Critical Reviews in Food Science and Nutrition*, 57(16), 3543-3564. <https://doi.org/10.1080/10408398.2016.1180501>
- [14] Canfora, E. E., Jocken, J. W. E., & Blaak, E. E. (2015). Short-chain fatty acids in control of body weight and insulin sensitivity. *Nature Reviews Endocrinology*, 11(10), 577-591. <https://doi.org/10.1038/nrendo.2015.128>
- [15] Chambers, L., McCrickerd, K., & Yeomans, M. R. (2015). Optimizing foods for satiety. *Trends in Food Science & Technology*, 41(2), 149-160. <https://doi.org/10.1016/j.tifs.2014.10.007>
- [16] Mozaffarian, D., Griffin, B. A., Manson, J. E., et al. (2021). Role of nutritional science in clinical medicine: A review. *BMJ*, 372, n209. <https://doi.org/10.1136/bmj.n209>
- [17] United States Department of Agriculture (USDA). (2023). FoodData Central: Wheat flour, refined. Retrieved from <https://fdc.nal.usda.gov/>
- [18] FatSecret. (2024). Whole wheat flour nutrition facts. Retrieved from <https://www.fatsecret.com/>
- [19] European Food Information Resource (EuroFIR). (2022). Food composition tables for cereals and derived products.
- [20] Doza Zilnică Recomandată (DZR). Romedic.ro. Available online: Romedic - Doza Zilnică Recomandată (accessed on 1 May 2026).

- [21] European Food Safety Authority Dietary Reference Values for Nutrients. EFSA. Available online: EFSA Dietary Reference Values (accessed on 1 May 2026).
- [22] World Health Organization Healthy Diet Fact Sheet. WHO. Available online: WHO Healthy Diet Fact Sheet (accessed on 1 May 2026).
- [23] Fardet, A. New hypotheses for the health-protective mechanisms of whole-grain cereals: What is beyond fibre? *Nutr. Res. Rev.* 2010, 23, 65-134. <https://doi.org/10.1017/S0954422410000041>.
- [24] Slavin, J. Fiber and Prebiotics: Mechanisms and Health Benefits. *Nutrients* 2013, 5, 1417-1435. <https://doi.org/10.3390/nu5041417>.
- [25] Makki, K.; Deehan, E.C.; Walter, J.; Bäckhed, F. The Impact of Dietary Fiber on Gut Microbiota in Host Health and Disease. *Cell Host Microbe* 2018, 23, 705-715. <https://doi.org/10.1016/j.chom.2018.05.012>.
- [26] Clark, M.J.; Slavin, J.L. The Effect of Fiber on Satiety and Food Intake: A Systematic Review. *J. Am. Coll. Nutr.* 2013, 32, 200-211. <https://doi.org/10.1080/07315724.2013.791194>.
- [27] World Health Organization Healthy Diet Fact Sheet. Available online: WHO Healthy Diet Fact Sheet (accessed on 13 May 2026)